

## **ELECTRONIC PRESCRIBING (E-PRESCRIBING)**

## Compounded Semaglutide Img-Iml in SubMagna™ HMW

- 1. Identify D&H Drugstore Broadway as the Desired Pharmacy in EMR System
  - Phone Number: 573-777-7333
  - Address: 1001 West Broadway Columbia, MO 65203
  - NPI: 1982690384
- 2. Select the medication Two Options Typically Considered
  - If your EMR system allows you to free-form a compounded medication that does not have a singlenumber NDC, consider free-form typing "Compounded Semaglutide Img-ImI in SubMagna™ HMW" as the prescribed medication.
  - If your EMR system does NOT allow you to free-form a compounded medication, consider selecting RYBELSUS® 14mg Tablet and adding the clarification for the specific medication prescribed in the SIG or Prescription Direction Field as described below in #3.
- 3. Add the Directions or "SIG" (Use to Embed Compound Name and Compound Directions Pick One)
  - Compounded Semaglutide Img-Iml in SubMagna<sup>™</sup> HMW Place 0.5ml under the tongue for minimum of 90 seconds before swallowing once daily for 1 week. Increase to Iml once daily for 2 to 4 weeks and then contact prescriber for dosage escalation if needed. Do not eat or drink for 30 minutes after taking medication.
  - Compounded Semaglutide 1mg-1ml in SubMagna<sup>™</sup> HMW Once daily, place \_\_\_\_ml under tongue for a minimum of 90 seconds before swallowing. Do not eat or drink for 30 minutes after taking.
- 4. Select Quantity in 30ml increments
- 5. Enter Number of Authorized Refills

RYBELSUS® is a registered trademark of Novo Nordisk A/S. No compounded medication is reviewed by FDA for safety or efficacy. SubMagna™ HMW is a trademark of Kingdom Licensing. Above is general proposed guidance that should be adapted to each individual prescriber's practice setting.

## **QUESTIONS? CONTACT US!**